Peloquin, P., see Townsend, G. (9) 122
Pérez-Martínez, L., see Joseph-Bravo, P. (9) 93
Pisu, M.G., see Trapani, G. (9) 139
Pubill, D., see Verdaguer, E. (9) 41

Rampin, O., see Ghanima, A. (9) 1
Reho, A., see Trapani, G. (9) 139
Rimbau, V., see Verdaguer, E. (9) 41
Ríos, C., see Montes, S. (9) 9
Rougier, A., see Leré, C. (9) 49

Schmid-Elsaesser, R., see Zausinger, S. (9) 112
Schweitzer, L., see Heck, W.L. (9) 173
Seu, E., see Trapani, G. (9) 139
Slusarczyk, A., see Heck, W.L. (9) 173
Smeyne, M. and Smeyne, R.J.
Method for culturing postnatal substantia nigra as an in vitro model of experimental Parkinson's disease (9) 105
Smeyne, R.J., see Smeyne, M. (9) 105
Smith, C.P., Somogyi, G.T., Bird, E.T., Chancellor, M.B. and Boone, T.B.
Neurogenic bladder model for spinal cord injury: spinal cord microdialysis and chronic urodynamics (9) 57
Smith, P.F., see Horii, A. (9) 77
Somogyi, G.T., see Smith, C.P. (9) 57
Suzuki, Y., see Zhao, B.-Q. (9) 85

Tang, H.M., see Ludvig, N. (9) 23
Terwilliger, R., see Newton, S.S. (9) 214
Townsend, G., Peloquin, P., Kloosterman, F., Hetke, J.F. and Leung, L.S.
Recording and marking with silicon multichannel electrodes (9) 122
Trapani, G., Dazzi, L., Pisu, M.G., Reho, A., Seu, E. and Biggio, G.
A rapid method for obtaining finasteride, a 5α-reductase inhibitor, from commercial tablets (9) 139
Tsuchiya, H., see Tsuji, S. (9) 16
Tsuji, S., Nakatomi, R., Tsuchiya, H., Hirai, K., Katayama, Y., Motelica-Heino, I. and Hashikawa, T.
Perineuronal surface acetylcholinesterase activity of a fine neural network stained histochemically and observed with backscattered electron imaging and X-ray mapping methods (9) 16

Umemura, K., see Zhao, B.-Q. (9) 85

Verdaguer, E., Pubill, D., Rimbau, V., Jiménez, A., Escubedo, E., Camarasa, J., Pallàs, M. and Camins, A.
Evaluation of neuronal cell death by laser scanning cytometry (9) 41

Wagner, G.C., see Michna, L. (9) 181
Wills, K.V., see Martí, J. (9) 197
Wrang, M.L., see Alsbo, C.W. (9) 157

Yang, X. and Marchand, J.E.
A sensitive technique to clone low abundance receptor transcripts from single microdissected tissue punches (9) 144

Zausinger, S., Baethmann, A. and Schmid-Elsaesser, R.
Anesthetic methods in rats determine outcome after experimental focal cerebral ischemia: mechanical ventilation is required to obtain controlled experimental conditions (9) 112
Zhao, B.-Q., Suzuki, Y., Kondo, K., Kawano, K.-i., Ikeda, Y. and Umemura, K.
A novel MCA occlusion model of photothrombotic ischemia with cyclic flow reductions: development of cerebral hemorrhage induced by heparin (9) 85